

POSTER PRESENTATION

Open Access

Novel adenoviral serotype 5 based immunotherapeutic induces T-cell responses despite anti-adenoviral neutralizing antibodies in colorectal cancer patients

Michael Morse², Elizabeth Gabitzsch^{1*}, Joseph P Balint Jr.¹, Younong Xu¹, HK Lyerly², Frank Jones¹

From Society for Immunotherapy of Cancer 28th Annual Meeting
National Harbor, MD, USA. 8-10 November 2013

Naturally occurring or induced Ad-specific neutralizing antibodies impeded the activity of recombinant Ad5-based vectors with E1 deletions. An improved Ad vector with deletions of the E1 and the E2b regions (Ad5 [E1-, E2b-]), the latter encoding the DNA polymerase and the pre-terminal protein, with significantly diminished late phase viral protein expression, were hypothesized to avoid immunological clearance and induce more potent immune responses against the encoded tumor antigen transgene in Ad-immune hosts. In the present phase I/II study, cohorts of patients with advanced metastatic colorectal cancer (mCRC) were immunized with escalating doses of Ad5 [E1-, E2b-]-CEA(6D). The carcinoembryonic antigen (CEA) transgene employed contains a modification (CAP1-6D) designed to enhance CTL stimulation. CEA-specific CMI responses were observed despite the presence of pre-existing Ad5 immunity in a majority (61.3%) of patients. Long-term follow-up in some of the patients revealed a waning of the induced CEA directed immune responses at around 6 months. Importantly, there was minimal toxicity, and overall patient survival (48% at 12 months) was similar regardless of pre-existing Ad5 neutralizing antibody titers. Our patient demographics, albeit limited in size, were similar with previously published studies of patients with chemotherapy-refractory mCRC. Of particular interest is the observation that treated mCRC patients in our study exhibited favorable survival probability. The results demonstrate that, in cancer

patients, the novel Ad5 [E1-, E2b-] gene delivery platform generates significant CMI responses to the tumor antigen CEA in the setting of both naturally acquired and immunization-induced Ad5-specific immunity.

Authors' details

¹Etubics Corporation, Seattle, WA, USA. ²Duke University Medical Center, Durham, NC, USA.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P129

Cite this article as: Morse et al.: Novel adenoviral serotype 5 based immunotherapeutic induces T-cell responses despite anti-adenoviral neutralizing antibodies in colorectal cancer patients. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P129.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Etubics Corporation, Seattle, WA, USA

Full list of author information is available at the end of the article