

POSTER PRESENTATION

Open Access

IL-12 primed CD8+ T-cells possess enhanced persistence and anti-tumor efficacy because of greater IL-7 responsiveness

C Bryce Johnson*, Bennett R May, Colleen A Cloud, Mark P Rubinstein, David J Cole

From Society for Immunotherapy of Cancer 28th Annual Meeting
National Harbor, MD, USA. 8-10 November 2013

Ex vivo IL-12 priming of tumor-reactive CD8+ T-cells enhances their persistence and subsequent anti-tumor efficacy upon adoptive cell transfer (ACT) into lymphodepleted mice. Since IL-7 and IL-15 are considered critical for the persistence of adoptively transferred T-cells, we investigated the responsiveness of IL-12 primed CD8+ T-cells to these homeostatic cytokines to help elucidate the mechanisms behind their superior anti-tumor abilities. Using pmel-1 T-cell receptor transgenic mice, we found that pmel CD8+ T cells activated in the presence of IL-12 (pmelIL-12) showed much greater expansion in an irradiated (6 Gy) host compared to cells activated without IL-12 (pmelsham). This expansion was dependent on host IL-7, but not IL-15. Compared to pmelsham, pmelIL-12 demonstrated greatly enhanced in vitro IL-7 responsiveness, as measured by proliferation and phosphorylation of signaling molecules STAT5, S6 and AKT. These striking differences were not seen with IL-15. Despite not playing a major role in pmelIL-12 effector phase expansion, IL-15 was critical for maximum anti-tumor efficacy of pmelIL-12, as irradiated mice devoid of IL-15, like mice receiving IL-7 neutralizing antibody (clone M25), showed reduced survival compared to irradiated mice receiving pmelIL-12 only. This decreased tumor control likely occurred because IL-15 was needed for long-term persistence of pmelIL-12. Together, these findings suggest that IL-12 priming of CD8+ T-cells augments the efficacy of ACT protocols in part by conferring an enhanced ability to respond to IL-7, thereby enabling transferred cells to persist in the IL-7 rich post-lymphodepletion environment.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P17

Cite this article as: Johnson *et al.*: IL-12 primed CD8+ T-cells possess enhanced persistence and anti-tumor efficacy because of greater IL-7 responsiveness. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P17.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Medical University of South Carolina, Charleston, SC, USA



© 2013 Johnson *et al.*; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.