POSTER PRESENTATION

Open Access

Open-label, randomized, multi-center study comparing the sequence of high dose Aldesleukin (Proleukin® (HD IL-2) and Ipilimumab Yervoy®) in patients with metastatic melanoma (proclivity 02)

Sapna Patel¹, Mohammed Milhem², Sigrun Hallmeyer³, Gregory Daniels⁴, Lee Cranmer⁵, Bret Taback⁶, Lawrence Flaherty⁷, Sandra Aung⁸, James Lowder^{8*}, William Sharfman⁹

From Society for Immunotherapy of Cancer 29th Annual Meeting
National Harbor, MD, USA. 6-9 November 2014

Purpose

To investigate whether the sequence of HD IL-2 and a checkpoint inhibitor, Ipilimumab, will have additive or synergistic efficacy or toxicity when used in rapid sequence.

Schema

Adult patients with Stage IV or unresectable Stage III metastatic melanoma who are eligible to receive HD IL-2, treatment naïve or have received prior adjuvant therapy are randomized to a sequential administration

⁸Prometheus Laboratories, San Diego, CA, USA
Full list of author information is available at the end of the article

of 4 doses of Ipilimumab or 4 cycles of HD IL-2 dosed according to their package inserts (figure 1). Fifty of the patients will start with one drug and 50 the other. The second drug will begin as soon as practically possible, without waiting for relapse. Entry criteria have recently been amended to include prior treatment with anti-PD-1 or anti-PDL-1. Twelve US sites are currently enrolling patients. An independent Data and Safety Monitoring Committee oversees the study. The primary endpoint is the proportional one year survival in the ITT population and a protocol defined population of patients who have received at least half of the planned doses of both study drugs. Clinical response and progression free survival will also be assessed. The primary endpoint is the proportional one year survival in the ITT population and a protocol defined population of patients who have received at least half of the planned doses of both study drugs. Clinical response and progression free survival will also be assessed.

Current status

Twelve US sites are currently enrolling patients. To date 16 patients have been enrolled, 9 on the Ipilimumab first and 7 on the HD IL-2 first arms. No synergistic toxicity has been observed, but one death occurred in the HD IL-2 arm and one colectomy on the Ipilimumab arm, both prior to administration of the other drug.

Authors' details

¹MD Anderson Cancer Center, Houston, TX, USA. ²University of Iowa, Iowa City, IA, USA. ³Oncology Specialists SC, Park Ridge, IL, USA. ⁴Moore's Cancer Center, La Jolla, CA, USA. ⁵University of Arizona Cancer Center, Tucson, AZ, USA. ⁶Columbia University, New York, NY, USA. ⁷Karmanos Cancer Center, Detroit, MI, USA. ⁸Prometheus Laboratories, San Diego, CA, USA. ⁹Johns Hopkins University, Baltimore, MD, USA.

Published: 6 November 2014

doi:10.1186/2051-1426-2-S3-P78

Cite this article as: Patel et al.: Open-label, randomized, multi-center study comparing the sequence of high dose Aldesleukin (Proleukin® (HD IL-2) and Ipilimumab Yervoy®) in patients with metastatic melanoma (proclivity 02). *Journal for ImmunoTherapy of Cancer* 2014 2 (Suppl 3):P78.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit