

POSTER PRESENTATION

Open Access

Integrated whole genome and transcriptome analysis identified a therapeutic minor histocompatibility antigen encoded by an alternative ITGB2 transcript

Margot J Pont^{1*}, Dyantha van der Lee¹, Edith van der Meijden¹, Cornelis van Bergen¹, Michel Kester¹, Maria Honders¹, Martijn Vermaat², Matthias Eefting¹, Erik Marijt¹, Szymon Kielbasa³, Peter-Bram 't Hoen², Frederik Falkenburg¹, Marieke Griffioen¹

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

Patients with hematological malignancies can be successfully treated with allogeneic stem cell transplantation (alloSCT). In HLA-matched alloSCT, donor T cells can mediate desired anti-tumor immunity as well as undesired side effects by recognizing minor histocompatibility antigens (MiHA) on patient cells. MiHA are polymorphic peptides with amino acid changes that are created by genetic variants and recognized by specific T cells. MiHA with hematopoiesis restricted expression are relevant targets for immunotherapy to augment anti-tumor immunity after alloSCT without side effects.

For high-throughput discovery of MiHA, we previously developed whole genome association scanning (WGAs) in which T cell recognition of a panel of EBV-B cell lines is investigated for association with single nucleotide polymorphisms (SNPs) to identify the genomic region that encodes the antigen. Although WGAs is an efficient strategy for MiHA discovery, strong association with SNPs in introns or other regions outside coding exons can be found for approximately one third of the T cell clones, whereas no SNP disparity in the primary transcript is present between patient and donor, suggesting that the antigen may be encoded by an alternative transcript. To identify MiHA encoded by alternative transcripts, we developed an integrated approach in which WGAs is combined with whole transcriptome data from the GEUVADIS project.

By performing WGAs, we identified associating SNPs for two HLA-B*15:01-restricted MiHA that were targeted by donor CD8 T cells in a patient with strong anti-tumor immunity after HLA-matched alloSCT. One antigen (LB-GLE1-1V) was encoded by an associating SNP in exon 6 of the *GLE1* gene. For the other antigen, an associating SNP in intron 3 of the ITGB2 gene was found, but no SNP disparity was present in the normal ITGB2 transcript between patient and donor. Therefore, we investigated whether the antigen may be encoded by an alternative transcript. Using RNA-sequence data, we identified an alternative ITGB2 transcript in which part of intron 3 was retained. Q-PCR analysis showed that expression of this transcript was restricted to hematopoietic cells from SNP-positive individuals. In silico translation of the transcript revealed a peptide with strong predicted binding to HLA-B*15:01 (LB-ITGB2-1), which was recognized by specific T cells. T cells for LB-ITGB2-1 also recognized and lysed leukemic cells of different origins, while no reactivity against patient fibroblasts could be detected.

In summary, RNA-sequence analysis enabled discovery of LB-ITGB2-1 as immune epitope encoded by an alternative gene transcript. Our data support the therapeutic relevance of LB-ITGB2-1 as target for T cell therapy to stimulate anti-tumor immunity after alloSCT.

¹Department of Hematology, Leiden University Medical Center, Leiden, Netherlands Authors' details

¹Department of Hematology, Leiden University Medical Center, Leiden, Netherlands. ²Department of Human Genetics, Leiden University Medical

Full list of author information is available at the end of the article



© 2015 Pont et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http:// creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/ zero/1.0/) applies to the data made available in this article, unless otherwise stated. Center, Leiden, Netherlands. ³Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, Leiden, Netherlands.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P285 Cite this article as: Pont *et al.*: Integrated whole genome and transcriptome analysis identified a therapeutic minor histocompatibility antigen encoded by an alternative ITGB2 transcript. *Journal for ImmunoTherapy of Cancer* 2015 3(Suppl 2):P285.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

BioMed Central

Submit your manuscript at www.biomedcentral.com/submit