

POSTER PRESENTATION

Open Access

Impact of low-lipid environment on de novo lipid synthesis pathways in cancer cells

Nousheen Zaidi

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

Increased lipogenesis is a hallmark of a wide variety of cancers and is under intense investigation as potential anti-neoplastic target. Although brisk lipogenesis is observed in the presence of exogenous lipids, evidence is mounting that these lipids may adversely affect the efficacy of inhibitors of lipogenic pathways. Therefore, to fully exploit the therapeutic potential of lipid synthesis inhibitors, a better understanding of the interrelationship between *de novo* lipid synthesis and exogenous lipids and their respective role in cancer cell proliferation and therapeutic response to lipogenesis inhibitors is of critical importance. Here, we show that the proliferation of various cancer cell lines (PC3M, HepG2, HOP62 and T24) is attenuated when cultured in lipid-reduced conditions in a cell line-dependent manner, with PC3M being the least affected. Interestingly, all cell lines - lipogenic (PC3M, HepG2, HOP62) as well as non-lipogenic (T24) - raised their lipogenic activity in these conditions, albeit to a different degree. Cells that attained the highest lipogenic activity under these conditions were best able to cope with lipid reduction in term of proliferative capacity. Supplementation of the medium with very low density lipoproteins, free fatty acids and cholesterol reversed this activation, indicating that the mere lack of lipids is sufficient to activate *de novo* lipogenesis in cancer cells. Consequently, cancer cells grown in lipid-reduced conditions became more dependent on *de novo* lipid synthesis pathways and were more sensitive to inhibitors of lipogenic pathways, like Soraphen A and Simvastatin. Collectively, these data indicate that limitation of access to exogenous lipids, as may occur in intact tumors, activates *de novo* lipogenesis in cancer cells, helps them to thrive under these conditions and makes them more vulnerable to lipogenesis inhibitors. These observations have important implications for the

design of new anti-neoplastic strategies targeting the cancer cell's lipid metabolism.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P424

Cite this article as: Zaidi: Impact of low-lipid environment on de novo lipid synthesis pathways in cancer cells. *Journal for ImmunoTherapy of Cancer* 2015 3(Suppl 2):P424.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



University of the Punjab, Lahore, Pakistan



© 2015 Zaidi This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.